

Solar Bulletin

THE AMERICAN ASSOCIATION OF VARIABLE STAR OBSERVERS
SOLAR SECTION



Rodney Howe, Editor, Chair
c/o AAVSO, 49 Bay State Rd
Cambridge, MA 02138 USA

Web: <http://www.aavso.org/solar-bulletin>
Email: solar@aavso.org
ISSN 0271-8480

Volume 74 Number 10

October 2018

The Solar Bulletin of the AAVSO is a summary of each month's solar activity recorded by visual solar observers' counts of group and sunspots, and the VLF radio recordings of SID Events in the ionosphere. Section 1 gives contributions by our members. The sudden ionospheric disturbance report is in Section 2. The relative sunspot numbers are in Section 3. Section 4 has endnotes.

1 Laboratory for Atmospheric and Space Physics (LASP)




Figure 1: LASP has a new web site, Lasp Interactive Solar IRadiance Datacenter (LISIRD) which houses solar data from many observatories. And now they display the AAVSO daily and monthly R_a back to 1944. (<http://lasp.colorado.edu/lisird/>)

For further reading on LASP: (<http://lasp.colorado.edu/home/>)

2 Sudden Ionospheric Disturbance (SID) Report

2.1 SID Records

October 2018 (Figure 2): There were two B class flares on the 12th of October recorded here Fort Collins, CO. However, there is only a slight possibility the B7.2 flare was recorded from NLK as an ionosphere SID event.




Figure 2: VLF recording at Fort Collins, Colorado.

2.2 SID Observers

In October 2018 we had 16 AAVSO SID observers who submitted VLF data as listed in Table 1. There were a few observers who recorded SID events on the 12th and 13th this month.

Table 1: 201810 VLF Observers

Observer	Code	Stations
A McWilliams	A94	NML
J Wallace	A97	NAA
L Loudet	A118	DHO
J Godet	A119	GBZ ICV
B Terrill	A120	NWC
F Adamson	A122	NWC
S Oatney	A125	NML NLK NAA
J Karlovsky	A131	NSY ICV
S Aguirre	A138	NPM
G Silvis	A141	NLK
I Ryumshin	A142	GQD DHO
R Rogge	A143	GQD
K Menzies	A146	NAA
R Russel	A147	NPM
L Ferreira	A149	NWC
A Maevsky	A151	DHO GQD ICV

Figure 3 depicts the importance rating of the solar events. The durations in minutes are -1: LT 19, 1: 19-25, 1+: 26-32, 2: 33-45, 2+: 46-85, 3: 86-125, and 3+: GT 125.




Figure 3: VLF SID Events.

2.3 Solar Flare Summary from GOES-15 Data

In October 2018, There were seven solar flare events. Five B class and 2 A class flares. A little more flaring this month compared to last month. There were 26 days this month with no GOES-15 reports of flares. (see Figure 4).




Figure 4: GOES - 15 XRA flares

3 Relative Sunspot Numbers R_a

Reporting monthly sunspot numbers consists of submitting an individual observer's daily counts for a specific month to the AAVSO Solar Section. These data are maintained in a SQL database. The monthly data then are extracted for analysis. This section is the portion of the analysis concerned with both the raw and daily average counts for a particular month. Scrubbing and filtering the data assure error-free data are used to determine the monthly sunspot numbers.

3.1 Raw Sunspot Counts

The raw daily sunspot counts consist of submitted counts from all observers who provided data in October 2018. These counts are reported by the day of the month, and are either from data not scrubbed or corrected data.

The reported raw daily average counts have been checked for errors and inconsistencies, and no known errors are present. All observers whose submissions qualify through this month's scrubbing process are represented in Figure 5.




Figure 5: Raw Wolf number average, minimum and maximum by day of the month for all observers.




Figure 6: Raw Wolf average and R_a numbers by day of the month for all observers.

3.2 American Relative Sunspot Numbers

The relative sunspot numbers, R_a , contain the sunspot numbers after the submitted data are scrubbed and modeled by Shapley's method with k -factors (<http://iopscience.iop.org/article/10.1086/126109/pdf>). The Shapley method is a statistical model that agglomerates variation due to random effects such as observer and fixed effects such as seeing condition. The raw Wolf averages and calculated R_a are seen in Figure 6 and Table 2 shows the Day (column 1) of the observation, the Number of Observations is in column 2, the raw Wolf number is in column 3, and the Shapley correction (R_a) is in column 4.

Table 2: 201810 American Relative Sunspot Numbers (R_a).

Day	Number of Observers	Raw	R_a
1	32	16	11
2	33	13	8
3	36	13	8
4	30	6	4
5	34	0	0
6	30	0	0
7	30	0	0
8	28	0	0
9	32	0	0
10	28	0	0
11	26	4	3
12	37	13	9
13	30	16	12
14	34	14	9
15	26	8	6

Continued

Table 2: 201810 American Relative Sunspot Numbers (R_a).

Day	Number of		
	Observers	Raw	R_a
16	35	0	0
17	30	4	3
18	36	0	0
19	36	0	0
20	44	0	0
21	46	4	3
22	41	0	0
23	32	0	0
24	34	0	0
25	27	0	0
26	28	0	0
27	29	0	0
28	31	0	0
29	31	0	0
30	34	0	0
31	30	0	0
Averages	32.6	3.6	2.5

3.3 Sunspot Observers

Table 3 lists the Observer Code (column 1), the Number of Observations (column 2) submitted for October 2018, and the Observer Name (column 3). The final rows of the table give the total number of observers who submitted sunspot counts and the total number of observations submitted. The total number of observers is 65 and the total number of observations is 1010.

Table 3: 201810 Number of observations by observer

Observer Code	Number of Observations	Observer Name
AAP	2	A. Patrick Abbott
AAX	12	Alexandre Amorim
AJV	20	J. Alonso
ARAG	30	Gema Araujo
ASA	24	Salvador Aguirre
ATE	4	Teofilo Arranz Heras
BARH	16	Howard Barnes
BATR	4	Roberto Battaiola
BERJ	23	Jose Alberto Berdejo
BLAJ	3	John A. Blackwell
BMF	17	Michael Boschat
BRAD	25	David Branchett
BRAF	6	Raffaello Braga
BROB	29	Robert Brown

Continued

Table 3: 201810 Number of observations by observer

Observer Code	Number of Observers	Observer Name
BSAB	25	Santanu Basu
CHAG	27	German Morales Chavez
CIOA	24	Ioannis Chouinavas
CKB	19	Brian Cudnik
CNT	21	Dean Chantiles
DEMF	5	Frank Dempsey
DJOB	5	Jorge del Rosario
DMIB	16	Michel Deconinck
DROB	4	Bob Dudley
DUBF	23	Franky Dubois
EHOA	19	Howard Eskildsen
ERB	17	Bob Eramia
FERJ	20	Javier Ruiz Fernandez
FLET	19	Tom Fleming
FLF	4	Fredirico Luiz Funari
FTAA	6	Tadeusz Figiel
FUJK	23	K. Fujimori
HAYK	7	Kim Hay
HOWR	22	Rodney Howe
JDAC	8	David Jackson
JENS	3	Simon Jenner
JGE	1	Gerardo Jimenez Lopez
KAND	24	Kandilli Observatory
KAPJ	15	John Kaplan
KNJS	31	James & Shirley Knight
KROL	27	Larry Krozel
LEVM	16	Monty Leventhal
LKR	3	Kristine Larsen
LRRA	9	Robert Little
MCE	22	Etsuiku Mochizuki
MILJ	19	Jay Miller
MJAF	30	Juan Antonio Moreno Quesada
MJHA	21	John McCammon
MUDG	8	George Mudry
MWU	10	Walter Maluf
ONJ	11	John O'Neill
RLM	10	Mat Raymonde
SDOH	31	Solar Dynamics Obs - HMI
SMNA	6	Michael Stephanou
SNE	3	Neil Simmons
SONA	18	Andries Son
STAB	25	Brian Gordon-States
SUZM	21	Miyoshi Suzuki

Continued

Table 3: 201810 Number of observations by observer

Observer Code	Number of Observers	Observer Name
TESD	25	David Teske
TPJB	5	Patrick Thibault
TST	6	Steven Toothman
URBP	21	Piotr Urbanski
VARG	22	A. Gonzalo Vargas
VIDD	12	Daniel Vidican
WGI	4	Guido Wollenhaupt
WILW	22	William M. Wilson
Totals	1010	65

3.4 Generalized Linear Model of Sunspot Numbers

Dr. Jamie Riggs, Solar System Science Section Head, International Astrostatistics Association, maintains a relative sunspot number (R_a) model containing the sunspot numbers after the submitted data are scrubbed and modeled by a Generalized Linear Mixed Model (GLMM), which is a different model method from the Shapley method of calculating R_a in Section 3 above. The GLMM is a statistical model that accounts for variation due to random effects and fixed effects. For the GLMM R_a model random effects include the AAVSO observer as these observers are a selection from all possible observers, and the fixed effects include seeing conditions at one of four possible levels. More details on GLMM are available in a paper (GLMM05) on <http://www.spesi.org/?page.id=65> of the sunspot counts research page. The paper title is *A Generalized Linear Mixed Model for Enumerated Sunspots*.

Figure 7 shows the monthly GLMM R_a numbers for the 24th solar cycle to date. The solid cyan curve that connects the red X's is the GLMM model R_a estimates of excellent seeing conditions, which in part explains why these R_a estimates often are higher than the Shapley R_a values. The dotted black curves on either side of the cyan curve depict a 99% confidence band about the GLMM estimates. The confidence band uses the large sample approximation based on the Gaussian distribution. The green dotted curve connecting the green triangles is the Shapley method R_a numbers. The dashed blue curve connecting the blue O's is the SILSO values for the monthly sunspot numbers.

The tan box plots for each month are the actual observations submitted by the AAVSO observers. The heavy solid lines approximately midway in the boxes represent the count medians. The box plot represents the InterQuartile Range (IQR), which depicts from the 25th through the 75th quartiles. The lower and upper whiskers extend 1.5 times the IQR below the 25th quartile, and 1.5 times the IQR above the 75th quartile. The black dots below and above the whiskers traditionally are considered outliers, but with GLMM modeling, they are observations that are accounted for by the GLMM model.

4 Endnotes

- Sunspot Reports: Kim Hay solar@aavso.org
- SID Solar Flare Reports: Rodney Howe ahowe@frii.com




Figure 7: GLMM fitted data for R_a . AAVSO data: <https://www.aavso.org/category/tags/solar-bulletin>. SILSO data: WDC-SILSO, Royal Observatory of Belgium, Brussels